

PLEASE RETURN:
COMPLETED ROSTER
COMPLETED TESTS
COMPLETED EVALUATION

PEGCO Inc.
532 N. RIDGEWOOD AVE
DAYTONA BEACH, FL. 32114
Phone: (386) 756-4266 Fax (386) 492-7821
WEB SITE: www.VolusiaCPR.com EMAIL: Bill@VolusiaCPR.com
NURSING HOME #NH 2772
ASSISTED LIVING #ALF 909
HOME HEALTH CARE #HH 1175
William Stacy RN

ROSTER

Program Title: **FALLS IN THE ELDERLY (1 HR)**

Date: _____ Facility _____
Name: _____

********PLEASE PRINT CLEARLY********

CERTIFICATES CAN NOW BE EMAILED IF WE CAN READ THE EMAIL
ADDRESS

1. NAME _____ PHONE _____
ADDRESS _____

EMAIL _____

2. NAME _____ PHONE _____
ADDRESS _____

EMAIL _____

3. NAME _____ PHONE _____
ADDRESS _____

EMAIL _____

4. NAME _____ PHONE _____
ADDRESS _____

EMAIL _____


FALLS IN THE ELDERLY

AGENDA

- 1. Introduction**
- 2. Geriatric Essentials**
- 3. Etiology**
- 4. Intrinsic factors**
- 5. Extrinsic factors**
- 6. Situational factors**
- 7. Complications**
- 8. Diagnosis**
- 9. History and physical examination**
- 10. Performance test**
- 11. Laboratory testing**
- 12. Prevention**

OBJECTIVES

- 1. Understand how important it is to ask about falls at least once per year.**
- 2. Identify the percentage of elderly people who will annually fall.**
- 3. List at three intrinsic factors related falls.**
- 4. Give two examples of extrinsic factors.**
- 5. Give two examples of situational factors.**
- 6. Identify some of the long term complications from falls.**
- 7. In diagnosing patients who maybe prone to falls, identify the test that can be used.**
- 8. Discuss the importance of a complete history and physical examination.**
- 9. List various laboratory test determined to be essential after the history and examination.**
- 10. Discuss the importance to focus on preventing or reducing the number of future falls and complications.**

[HOME](#) | [ABOUT MERCK](#) | [PRODUCTS](#) | [NEWSROOM](#) | [INVESTOR INFORMATION](#) | [CAREERS](#) | [RESEARCH](#) | [LICENSING](#) | [THE](#)

The Merck Manual of Geriatrics

[Contents](#)[Table Page](#)[Search the Book](#)[Index](#)

Section 2. Falls, Fractures, and Injury

Chapter 20. Falls

Topic: Falls

Falls

A fall results in a person coming to rest on the ground or another lower level; sometimes a body part strikes against an object that breaks the fall. Typically, events caused by acute disorders (eg, stroke, seizure) or overwhelming environmental hazards (eg, being struck by a moving object) are not considered falls. Most falls cause an injury; most injuries are minor, but some (eg, fractures, lacerations, head trauma) are serious. Falls most commonly occur when several different problems (eg, unsteady gait, imbalance, lower-extremity weakness, slower reflexes, environmental hazards) interact to overcome an elderly person's ability to compensate. After treatment of acute injuries, assessment aims to identify risk factors and appropriate interventions, thus decreasing the risk of future falls and fall-related injuries.

Geriatric Essentials

- Elderly patients should be asked about falls at least once per year because many patients do not volunteer that information.

About ½ of elderly people who fall cannot get back up without help, increasing their risk of dehydration, pressure ulcers, rhabdomyolysis, hypothermia, and pneumonia. Lower-extremity muscle strengthening and balance training reduce risk of falls in

elderly people.

Falls threaten the independence of elderly people and produce a cascade of individual and socioeconomic consequences. However, physicians are often unaware of falls because a routine history and physical examination typically does not include a specific evaluation for falls and because many people who fall do not have an obvious injury. Many elderly people are reluctant to report a fall because the resulting injury was minimal, because they attribute falling to the aging process, or because they fear being subsequently restricted in their activities or being institutionalized.

Annually, 30 to 40% of elderly people living in the community fall; 50% of nursing home residents fall. In the US, falls are the leading cause of accidental death and the 7th leading cause of death in people ≥ 65 ; 75% of deaths caused by falls occur in the 12.5% of the population that are ≥ 65 . In 1994, health care costs of fall-related injuries and complications were \$20 billion; in 2020, they are projected to reach \$44 billion.

Etiology

The best predictor of falling is a previous fall. However, falls in elderly people rarely have a single cause or risk factor. A fall is usually caused by a complex interaction among intrinsic factors (age-related decline in function, disorders, and adverse drug effects) extrinsic factors (environmental hazards), and situational factors (related to the activity being done--eg, rushing to the bathroom).

Intrinsic factors: Age-related changes can impair systems involved in maintaining balance and stability (eg, while standing, walking, or sitting). In the vestibular system, labyrinthine hair cells and nerve fibers are lost, preventing the brain from receiving vestibular sensory input. Proprioception in the lower extremities declines. Visual acuity, contrast sensitivity, depth perception, and dark adaptation decline. Changes in muscle activation patterns may impair the ability to maintain or recover balance in response to perturbations (eg, stepping onto an uneven surface, being bumped).

Chronic and acute disorders (see Table 20-1) and use of drugs (see Table 20-2) are major risk factors for falls. The risk of falls increase with the number of drugs taken. Psychoactive drugs are the drugs most commonly reported as increasing the risk of falls and fall-related injuries.

Extrinsic factors: Environmental factors can increase the risk of falls independently or, more importantly, by interacting with intrinsic factors (see Table 20-3). Risk is highest when the environment requires greater postural control and mobility (eg, when walking on a slippery surface) and when the environment is unfamiliar (eg, when relocated to a new home).

Situational factors: Certain activities or decisions may increase the risk of falls and fall-related injuries. Examples are walking in stocking feet or in footwear with high heels, rushing to the bathroom (especially at night when not fully awake or when lighting may be inadequate), and rushing to answer the telephone.

Complications

Falling, particularly falling repeatedly, increases risk of injury, hospitalization, and death. Longer-term complications can include decreased physical function, fear of falling, and institutionalization.

Over 50% of falls among elderly people result in an injury. Although most injuries are not serious (eg, affect soft tissue), fall-related injuries account for about 5% of hospitalizations in patients ≥ 65 . About 5% of falls result in fractures of the humerus, wrist, or pelvis. About 2% of falls result in a hip fracture. Other serious injuries (eg, head and internal injuries, lacerations) occur in about 10% of falls. Some fall-related injuries are fatal. About 5% of elderly people with hip fractures die while hospitalized; overall mortality in the 12 mo after a hip fracture ranges from 18 to 33%.

Many elderly people who fall are frail and have preexisting deficits in activities of daily living and instrumental activities of daily living these people are at increased risk of other complications after a fall. About $\frac{1}{2}$ cannot get up without help. Remaining on the floor for > 2

h after a fall increases risk of dehydration, pressure ulcers, rhabdomyolysis, hypothermia, and pneumonia.

Function and quality of life may deteriorate drastically after a fall; at least 50% of elderly people who were ambulatory before fracturing a hip do not recover their previous level of mobility. After falling, elderly people may fear falling again, so mobility is sometimes reduced because confidence is lost. Some people may even avoid certain activities (eg, shopping, cleaning) because of this fear. Decreased activity can increase joint stiffness and weakness, further reducing mobility. Falls reportedly contribute to 40% of nursing home admissions.

Diagnosis

Some falls are promptly recognized because of an obvious fall-related injury or concern about a possible injury. However, because elderly people often do not report falls, they should be asked about them at least once per year.

Patients who report a single fall should be evaluated for a balance or gait problem using the Get-Up-and-Go Test. For the test, patients are observed as they rise from a standard armchair, walk a fixed distance in a straight line, turn, walk back to the chair, and sit back down. Observation may detect lower-extremity weakness, imbalance while standing or sitting, or an unsteady gait.

Patients who require a more complete assessment of risk factors for falls include those who have difficulty during the Get-Up-and-Go Test, those who report multiple falls during screening, and those who are being evaluated after a recent fall (after acute injuries are identified and treated).

History and physical examination: When a more complete assessment of risk factors is needed, the focus is on identifying intrinsic, extrinsic, and situational factors that can be reduced by interventions targeted at them.

Patients are asked open-ended questions about the most recent fall o

falls, followed by more specific questions about when and where a fall occurred and what they were doing. Witnesses are asked the same questions. Patients should be asked whether they had premonitory or associated symptoms (eg, palpitations, shortness of breath, chest pain, vertigo, light-headedness) and whether consciousness was lost. Patients should also be asked whether any obvious extrinsic or situational factors may have been involved. The history should include questions about past and present medical problems, use of prescription and OTC drugs, and use of alcohol. Because eliminating all risk of future falls may be impossible, patients should be asked whether they could get back up without help after falling and whether any injuries occurred; the goal is reducing the risk of complications due to future falls.

The physical examination should be comprehensive enough to exclude obvious intrinsic causes of falls. If the fall occurred recently, temperature should be measured to determine whether fever was a factor. Heart rate and rhythm should be assessed to identify obvious bradycardia, resting tachycardia, or irregular rhythms. BP should be measured with patients supine and after patients stand for 1 and 5 min to rule out orthostatic hypotension. Auscultation can detect many types of valvular heart disorders. Visual acuity should be evaluated with patients wearing their usual corrective lenses if needed. Abnormalities in visual acuity should trigger a more detailed visual examination by an optometrist or ophthalmologist. The neck, spine, and extremities (especially the legs) should be evaluated for abnormalities, pain, and limitation in range of motion.

A neurologic examination should be done; it includes testing muscle strength and tone, sensation (including proprioception), coordination (including cerebellar function), stationary balance, and gait. Basic postural control and the proprioceptive and vestibular systems are evaluated using the Romberg test (in which patients stand with feet together and eyes closed). Tests to establish high-level balance function include the one-legged stance and tandem gait. If patients can stand on one leg for 10 sec with the eyes open and have an accurate 10-ft tandem gait, any intrinsic postural control deficit is

likely to be minimal. Vestibular function and mental status should be evaluated.

Performance tests: The Performance-Oriented Assessment of Mobility or Get-Up-and-Go Test can identify problems with balance and stability during walking and other movements that may indicate increased risk of falls.

Laboratory testing: There is no standard diagnostic evaluation. Testing should be based on the history and examination and helps rule out various causes: a CBC for anemia, plasma glucose measurement for hypoglycemia or hyperglycemia, and electrolyte measurement for dehydration. Tests such as ECG, ambulatory cardiac monitoring, and echocardiography are recommended only when a cardiac cause is suspected. Carotid massage under controlled conditions (IV access and cardiac monitoring) has been proposed to determine carotid hypersensitivity and ultimately who might respond to pacemaker treatment. The most appropriate use of carotid massage may be in patients with recurrent, unexplained falls, especially when the history involves falls of the abrupt "drop attack" variety. Spinal x-rays and cranial CT or MRI are indicated only when the history and physical examination detect new neurologic abnormalities.

Prevention

The focus is on preventing or reducing the number of future falls and fall-related injuries and complications, while maintaining as much of patients' function and independence as possible. For more information, see the Cochrane Library: Interventions for preventing falls, American Geriatrics Society: Guideline for the prevention of falls, and British Medical Journal: Interventions for the prevention of falls.

Patients who report a single fall and who do not have problems with balance or gait on the Get-Up-and-Go Test or a similar test should be given general information about reducing risk of falls. It should include how to use drugs safely and reduce environmental hazards (see Table 20-3).

Patients who have fallen more than once or who have problems during initial balance and gait testing should be referred to physical therapy or an exercise program. Physical therapy and exercise programs can be done in the home if patients have limited mobility. Physical therapists customize exercise programs to improve balance and gait and to correct specific problems contributing to fall risk. Some patients benefit from use of an assistive device (eg, cane, walker). Canes may be adequate for those with minimal unilateral weakness, but walkers, especially 4-wheel walkers, are more appropriate for patients with increased risk of falls attributable to bilateral leg weakness or impaired coordination. Physical therapists can help fit or size the devices and teach patients how to use them. More general exercise programs in health care or community settings can also improve balance and gait. For example, tai chi may be effective and can be done alone or in groups. Lower-extremity strength training helps many patients, but at least initially, such training should be supervised by a qualified therapist or trainer experienced in working with elderly people.

Drugs that can increase the risk of falls should be stopped, or their dosage should be adjusted to the lowest effective dose. If a specific disorder is identified as a risk factor, targeted interventions are required. For example, drugs and physical therapy may reduce risk for patients with Parkinson's disease. Vitamin D, particularly taken with Ca, can reduce fall risk; this reduction is greater than the accompanying improvement in musculoskeletal function. Pain management, physical therapy, and sometimes joint replacement surgery may reduce risk for patients with arthritis. A change to appropriate lenses (single lenses rather than bifocals or trifocals) or surgery helps some patients with visual impairment.

Correcting environmental hazards in the home may reduce the risk of falls (see Table 20-3). Elderly people should also be advised on how to reduce risk due to situational factors. For example, footwear with flat heels and firm midsoles is a better choice than footwear with high heels. Many people with chronic limited mobility (eg, severe arthritis, paresis) benefit from combined medical, rehabilitative, and environmental strategies. Wheelchair adaptations

(eg, removable foot plates to reduce tripping during transfers, antitip bars to prevent backward tipping), removable belts, and wedge seating may prevent falls in people with poor sitting balance or severe weakness when they are sitting or transferring. Restraints should not be used. Surveillance by a caregiver is more effective and safer. Motion detectors may be used, but a caregiver must be present to respond to the triggered alarm.

Hip protectors (padding sewn into special undergarments) may help protect elderly people who have fallen and are at risk of a hip injury, but many people are reluctant to wear protectors indefinitely. Compliant flooring (eg, firm rubber) can help dissipate the impact force, but a floor that is too compliant (eg, soft foam) may destabilize patients.

Patients who have fallen repeatedly should be evaluated for osteoporosis and, if osteoporosis is diagnosed, treated to reduce risk of fractures from any future falls. These patients should also be taught what to do if they fall and cannot get up. Useful techniques include turning from the supine position to the prone position, getting on all fours, crawling to a strong support surface, and pulling up. Having frequent contact with family members or friends, a phone that can be reached from the floor, a remote alarm, or a wearable emergency response system device can decrease the likelihood of lying on the floor for a long time after a fall.

This topic was last updated May 2005.

[CONTACT MERCK](#) | [SITE MAP](#) |  [PRIVACY POLICY](#) | [TERMS OF USE](#) | [COPYRIGHT © 1995-2007 MERCK & CO., INC.](#)

NAME _____ DATE _____

FALLS IN THE ELDERLY QUIZ

1. How often should a physician ask an elderly person regarding falling episodes? _____ . page 1

2. What is the percentage of elderly people who will fall annually.? _____ . Page 2

3. List three intrinsic factors related to falls: _____

_____ . Page 2

4. Give two examples of extrinsic factors related to falls

_____ . Page 3

5. Give two examples of situational factors related to falls _____

_____ . Page 3

6. List at least four long term complications of falls _____

_____ . Page 3

7. What is the name of the test a physician can use to evaluate balance or gait problems. _____ .

Page 4

8. Give three examples of neurological tests that should be done: _____

_____ . Page 5

9. What laboratory blood work should be included in the exam?

_____ . Page 6

10. Give four examples of preventive measures to reduce the frequency of falls in the elderly. _____

_____ . Page 7

PEGCO Inc.
532 N. RIDGEWOOD AVE
DAYTONA BEACH, FL. 32114
Phone: (386) 756-4266 Fax (386) 492-7821
WEB SITE: www.VolusiaCPR.com EMAIL: Bill@VolusiaCPR.com
NURSING HOME #NH 2772
ASSISTED LIVING #ALF 909
HOME HEALTH CARE #HH 1175

PROGRAM EVALUATION

COURSE TITLE: FALLS IN THE ELDERLY 1 HR

DATE: _____ LOCATION: PEGCO, Inc.

Please evaluate by circling the appropriate rating:

5-Excellent 4-Above average 3-Average 2-Fair 1-Poor

- | | |
|---|-----------------------------|
| 1. Overall quality of the program | 5 4 3 2 1 |
| 2. Overall content of the program | |
| a. content can improve my ability to perform my job | 5 4 3 2 1 |
| b. content reflected knowledge level and needs of learner | 5 4 3 2 1 |
| c. the material was current | 5 4 3 2 1 |
| 3. Achieved stated objectives | |
| a. total number of objectives in program _____ | |
| b. circle the number of met objectives | 1 2 3 4 5 6 7 8 9 <u>10</u> |
| c. the test material reflected the objectives listed | 5 4 3 2 1 |
| 4. Overall organization of the program | |
| a. material was organized to facilitate learning | 5 4 3 2 1 |
| b. material covered was adequate and accurate | 5 4 3 2 1 |

What did you like best about the program?

Your suggestions for improving this program.

Any topic ideas for future in-service programs

THANK YOU FOR USING PEGCO INC. WE APPRECIATE YOU.